arXiv:0909.4037v1 [math.PR] 22 Sep 2009

RANDOM INDUCED SUBGRAPHS OF CAYLEY GRAPHS INDUCED BY TRANSPOSITIONS

EMMA Y. JIN AND CHRISTIAN M. REIDYS⋆

Abstract. In this paper we study random induced subgraphs of Cayley graphs of the symmetric group induced by an arbitrary minimal generating set of transpositions. A random induced subgraph of this Cayley graph is obtained by selecting permutations with independent probability, λ_n. Our main result is that for any minimal generating set of transpositions, for probabilities $\lambda_n = \frac{1+\epsilon_n}{n-1}$ where $n^{-\frac{1}{3}+\delta} \le \epsilon_n < 1$ and $\delta > 0$, a random induced subgraph has a.s. a unique largest component of size $\wp(\epsilon_n)\frac{1+\epsilon_n}{n-1}n!$, where $\wp(\epsilon_n)$ is the survival probability of a specific branching process.

1. Introduction

One central problem arising in parallel computing is to determine an optimal linkage of a given collection of processors. A particular class of processor linkages with point-to-point communication links are static interconnection networks. The latter are widely used for message-passing architectures. A static interconnection network can be represented as a graph. The binary n-cubes, Q_2^n, [1, 27] are a particularly well-studied class of interconnection networks [10, 12, 13, 32].

Akers *et al.* [2] recognized the deficiencies of n-cubes as models for interconnection networks and proposed an alternative: the Cayley graph of the permutation group induced by the $(n-1)$ star-transpositions $(1\,i)$, which we denote by $\Gamma(S_n, P_n)$. Pak [28] studied minimal decompositions of a particular permutation via star-transpositions and Irving *et al.* [20] extended his results. The star-graph $\Gamma(S_n, P_n)$ is in many aspects superior to n-cubes [1, 27]. Some properties of star-graphs

---

Date: September, 2009.

2000 *Mathematics Subject Classification.* 05A16.

Key words and phrases. random graph, permutation, transposition, giant component, vertex boundary.

studied in [16, 18, 19, 17, 21, 25] were cycle-embeddings and path-embeddings. Diameter and fault diameter of star-graphs were computed by Akers *et al.* [2, 24, 31] and Lin *et al.* [26] analyzed diagnosability. An alternative to n-cubes as interconnection networks are the bubble-sort graphs [3], studied by Tchuente [33]. The bubble-sort graph is the Cayley graph of the permutation group induced by all $n-1$ canonical transpositions $(i\ i+1)$, denoted by $\Gamma(S_n, B_n)$.

Recently, Araki [5] brought the attention to a generalization of star- and bubble-sort graphs, the Cayley graph generated by all transpositions. The latter has direct connections to a problem of interest in computational biology: the evolutionary distances between species based on their genome order in the Cayley graph of signed permutations generated by reversals. A reversal is a special permutation that acts by flipping the order as well as the signs of a segment of genes. Hannenhalli and Pevzner [29] presented an algorithm computing minimal number of reversals needed to transform one sequence of distinct genes into a given signed permutation. For distant genomes, however, it is well-known, that the true evolutionary distance is generally much greater than the shortest distance [7, 22]. In order to obtain a more realistic estimate of the true evolutionary distance, the expected reversal distance was shifted into focus. Its computation, however, has proved to be hard and motivated to consider models better suited for computation. Point in case is the work of Eriksen *et al.* [11], where the authors derive a closed formula for the expected transposition distance and subsequently show how to use it as an approximation of the expected reversal distance.

In this paper we study a subgraph of the Cayley graph generated by all transpositions, the Cayley graph $\Gamma(S_n, T_n)$, where T_n is a minimal generating set of transpositions. Setting $T_n = P_n$ and $T_n = B_n$ we can recover the star- and the bubble-sort graph as particular instances. We study structural properties of $\Gamma(S_n, T_n)$ in terms of the random graph obtained by selecting permutations with independent probability. The main result of this paper is

Theorem 1. *Let* $\lambda_n = \frac{1+\epsilon_n}{n-1}$*, where* $n^{-\frac{1}{3}+\delta} \le \epsilon_n < 1$ *and* $\delta > 0$*. Let* T_n *be a minimal generating set of transpositions and let* $\Gamma_n = \Gamma_n(\lambda_n)$ *denote the random induced subgraph of* $\Gamma(S_n, T_n)$*, obtained by independently selecting each permutation with probability* λ_n*. Then* Γ_n *has a.s. a unique giant component,* $C_n^{(1)}$*, whose size is given by*

$$|C_n^{(1)}| \sim (1 - o(1)) \cdot \wp(\epsilon_n) \cdot \frac{1+\epsilon_n}{n-1} \cdot n!, \tag{1.1}$$

where $\wp(\epsilon_n) = (1 + o(1))x$ *for* $\epsilon_n = \epsilon > 0$*,* $x > 0$ *being the unique root of* $e^{-(1+\epsilon)y} = 1 - y$ *and* $\wp(\epsilon_n) = (2 + o(1))\epsilon_n$ *for* $0 < \epsilon_n = o(1)$*.*



Figure 1. The evolution of the giant component in random induced subgraphs of $\Gamma(S_9, P_9)$. We display the relative size of the giant component $\frac{|C_n^{(1)}|}{|\Gamma_n|}$ as a function of $(1+\epsilon)/(n-1)$ as data-curve (blue) versus the growth predicted by Theorem 1 (red).

Similar to the case of random induced subgraphs of n-cubes [30], random induced subgraphs of $\Gamma(S_n, T_n)$ exhibit a giant component for very small vertex selection probabilities. This is insofar interesting as important relations, like for instance, the isoperimetric inequality [14], do not play a key role.

The paper is organized as follows: after establishing in Section 2 some basic properties of $\Gamma(S_n, T_n)$ and preparing all necessary background, we analyze in Section 3 vertices contained in polynomial size subcomponents. The strategy is similar to that in [30], where first a specific branching process is embedded (for its first $\lfloor \frac{1}{4} n^{\frac{2}{3}} \rfloor$ steps) into $\Gamma(S_n, T_n)$. It is its survival probability that provides a lower bound on the probability that a given vertex is contained in a subcomponent of arbitrary, polynomial size. In Section 4 we "sandwich" this bound by showing that there are many vertices in "small" components. Only here we use $\epsilon < 1$. In Section 5 we show that there are many vertex disjoint paths between certain splits of permutations. The a.s. existence of the giant component follows using the ideas of Ajtai *et al.* [1].

2. Some basic facts

Let S_n denote the symmetric group over $[n]$. We write a permutation $\pi \in S_n$ as an n-tuple $(x_1, \dots, x_n)$. Let $T_n \subset S_n$ be a minimal generating set of transpositions. Plainly, we have for $i < j$

$$(x_1, \dots, x_i, \dots, x_j, \dots, x_n)\,(i\,j) = (x_1, \dots, x_j, \dots, x_i, \dots, x_n). \tag{2.1}$$

We consider the Cayley graph $\Gamma(S_n, T_n)$, having vertex set S_n and edges (v, v') where $v^{-1}v' \in T_n$. For $v, v' \in S_n$, let $d(v, v')$ be the minimal number of T_n-transpositions by which v and v' differ. For $A \subset S_n$ we set $\mathsf{B}(A, j) = \{v \in S_n \mid \exists \alpha \in A;\, d(v, \alpha) \le j\}$ and $\mathsf{d}(A) = \{v \in S_n \setminus A \mid \exists \alpha \in A;\, d(v, \alpha) = 1\}$ and call $\mathsf{B}(A, j)$ and $\mathsf{d}(A)$ the ball of radius j around A and the vertex boundary of A in $\Gamma(S_n, T_n)$. If $A = \{\alpha\}$ we simply write $\mathsf{B}(\alpha, j)$. Let $D, E \subset S_n$, we call D ℓ-dense in E if $\mathsf{B}(\sigma, \ell) \cap D \neq \varnothing$ for any $\sigma \in E$. Let "$\le$" be the following linear order over $\Gamma(S_n, T_n)$

$$\sigma \le \tau \quad\Longleftrightarrow\quad (d(\sigma, id) < d(\tau, id)) \;\vee\; (d(\sigma, id) = d(\tau, id) \;\wedge\; \sigma <_{\rm lex} \tau), \tag{2.2}$$

where id is the identity permutation and $<_{\rm lex}$ denotes the lexicographical order. Any notion of minimal or smallest element in a subset $A \in S_n$ refers to the linear order $\le$ of eq. (2.2).

Let $\Gamma_{\lambda_n}(S_n, T_n)$ be the probability space (random graph) consisting of $\Gamma(S_n, T_n)$-subgraphs, Γ_n, induced by selecting each $\Gamma(S_n, T_n)$-vertex with independent probability λ_n. A property M is a subset of induced subgraphs of $\Gamma(S_n, T_n)$ closed under graph isomorphisms. The terminology "M holds a.s." is equivalent to $\lim_{n\to\infty} \mathsf{Prob}(\mathsf{M}) = 1$. A component of Γ_n is a maximal, connected, induced Γ_n-subgraph, C_n. The largest Γ_n-component is denoted by $C_n^{(1)}$. We write $x_n \sim y_n$ if and only if (a) $\lim_{n\to\infty} x_n/y_n$ exists and (b) $\lim_{n\to\infty} x_n/y_n = 1$. We furthermore write $g(n) = O(f(n))$ and $g(n) = o(f(n))$ for $g(n)/f(n) \to \kappa$ as $n \to \infty$ and $g(n)/f(n) \to 0$ as $n \to \infty$, respectively. A largest component is called giant if it is unique in "size", i.e. any other component, C_n, satisfies $|C_n| = o(|C_n^{(1)}|)$.

Let $Z_n = \sum_{i=1}^n \xi_i$ be a sum of mutually independent indicator random variables (r.v.), ξ_i having values in $\{0, 1\}$. Then we have, [9], for $\eta > 0$ and $c_\eta = \min\{-\ln(e^\eta[1+\eta]^{-[1+\eta]}), \frac{\eta^2}{2}\}$

$$\mathsf{Prob}(\,|\,Z_n - \mathbb{E}[Z_n]\,|\, > \eta\, \mathbb{E}[Z_n]\,) \le 2e^{-c_\eta \mathbb{E}[Z_n]}\,. \tag{2.3}$$

n is always assumed to be sufficiently large and ϵ is a positive constant satisfying $0 < \epsilon < 1$. We write $B_m(\ell, \lambda_n) = \binom{m}{\ell}\lambda_n^\ell\,(1-\lambda_n)^{m-\ell}$.

Let us next recall some basic facts about branching processes, $\mathcal{P}_m = \mathcal{P}_m(p)$ [15, 23]. Suppose $\mathcal{P}_m$ is initiated at ξ. Let $(\xi_i^{(t)})$, $i, t \in \mathbb{N}$ count the number of "offspring" of the ith-"individual" of "generation" $(t-1)$, where the r.v. ξ and $\xi_i^{(t)}$ are $B_m(\ell, p)$-distributed. Let $\mathcal{P}_0 = \mathcal{P}_0(p)$ denote the branching process for which ξ is $B_m(\ell, p)$- and all $\xi_i^{(t)}$ are $B_{m-1}(\ell, p)$-distributed. Furthermore, let $\mathcal{P}_P(\lambda)$, $(\lambda > 0)$ denote the branching process in which the individuals generate offsprings according to the poisson distribution, i.e., $\mathbb{P}(\xi_i^{(t)} = j) = \frac{\lambda^j}{j!}e^{-\lambda}$. We consider the family of r.v. $(Z_i)_{i \in \mathbb{N}_0}$: $Z_0 = 1$ and $Z_t = \sum_{i=1}^{Z_{t-1}} \xi_i^{(t)}$ for $t \geq 1$ and interpret Z_t as the number of individuals "alive" in generation t. Of particular interest for us will be the limit $\lim_{t\to\infty} \mathbb{P}(Z_t > 0)$, i.e. the probability of infinite survival. We write

$$\pi_0(p) = \lim_{t\to\infty} \mathbb{P}_0(Z_t > 0),\ \pi_m(p) = \lim_{t\to\infty} \mathbb{P}_m(Z_t > 0) \text{ and } \pi_P(\lambda) = \lim_{t\to\infty} \mathbb{P}_P(Z_t > 0)$$

for the survival probability of $\mathcal{P}_0$, $\mathcal{P}_m$ and $\mathcal{P}_P(\lambda)$, respectively.

Corollary 1. [8] *Let $p = \lambda/n$.*
(1) If $\lambda > 1$ is fixed, then $\pi_0(p) = (1 + o(1))\pi_P(\lambda)$.
(2) Let $\lambda_n = 1 + \epsilon_n$, where $0 < \epsilon_n = o(1)$. Then, if $r = n - s$ and $s = o(n\epsilon_n)$,

$$\pi_0(p) = (1 + o(1))\pi_r(p) = (2 + o(1))\epsilon_n.$$

3. Components of size at least $O(n^{\frac{2}{3}+k\delta})$

Suppose $x > 0$ is the unique root of $e^{-(1+\epsilon)y} = 1 - y$ and

$$\wp(\epsilon_n) = \begin{cases} (1 + o(1))x & \text{for } \epsilon_n = \epsilon > 0 \\ (2 + o(1))\epsilon_n & \text{for } 0 < \epsilon_n = o(1). \end{cases} \tag{3.1}$$

For $k \in \mathbb{N}$ we set

$$\mu_n = \lfloor \frac{1}{2k(k+1)} n^{\frac{2}{3}} \rfloor, \quad \ell_n = \lfloor \frac{k}{2(k+1)} n^{\frac{2}{3}} \rfloor, \quad \text{and} \quad z_n = n - k\mu_n - \ell_n. \tag{3.2}$$

We next establish some basic properties of the Cayley graph $\Gamma(S_n, T_n)$:

Lemma 1. *Let T_n be a minimal generating set of S_n consisting of transpositions, then we have*
(1) T_n has cardinality $n - 1$ and corresponds uniquely to a labeled tree, $\mathcal{T}_n$ over $[n]$.
(2) there exists a sequence $(v_i)_{2\leq i}$ such that $T_n = \{(v_i\, s_i) \mid 2 \leq i \leq n\}$ and

$$\forall\, j < i; \quad ((x_1, \dots, x_n)(v_j\, s_j))_{v_i} \quad \neq \quad ((x_1, \dots, x_n)(v_i\, s_i))_{v_i}. \tag{3.3}$$

(3) *the diameter of* $\Gamma(S_n, T_n)$ *is given by*

$$\operatorname{diam}(\Gamma(S_n,T_n)) \le \sum_{i=2}^{n} \operatorname{diam}(\mathcal{T}_i) \le \binom{n}{2}. \tag{3.4}$$

Proof. It is straightforward to prove by induction that $|T_n| = n-1$. We next consider the graph $\mathcal{T}_n$ over $[n]$, having edge-set T_n. Since $\langle T_n\rangle = S_n$, $\mathcal{T}_n$ is connected and since T_n is independent, $\mathcal{T}_n$ is a tree. This establishes the mapping

$$\psi\colon \{T_n \mid T_n \text{ is a maximal independent transposition set}\} \longrightarrow \{\mathcal{T}_n \mid \mathcal{T}_n \text{ is a tree over } [n]\}.$$

Furthermore, ψ has an inverse; as the edges of a tree over $[n]$ give rise to a maximal independent set of transpositions that generate S_n, whence assertion (1).
In order to prove (2), we generate the tree $\mathcal{T}_n$ inductively as follows: we start with vertex 1 and consider its unique minimal element in distance 1, v_1, and set $\mathcal{T}_1 = \{(v_1\,1)\}$. Given $\mathcal{T}_i$, we consider the transposition $(v_{i+1}\,s_{i+1})$, where v_{i+1} is the unique minimal element contained in $\mathcal{T}_n \setminus \mathcal{T}_i$, having minimal distance to 1, and s_{i+1} is its unique $\mathcal{T}_i$-neighbor. We then set $\mathcal{T}_{i+1} = \mathcal{T}_i \cup \{(v_{i+1}\,s_{i+1})\}$. This process gives rise to the sequence of trees $\mathcal{T}_2 \subset \mathcal{T}_3 \subset \cdots \subset \mathcal{T}_n$ and denoting the vertex sets of $\mathcal{T}_i$ by V_i, we have $V_1 = \{1\} \subset V_2 \subset V_3 \subset \dots V_{n-1} \subset V_n = [n]$ where $\{v_i\} = V_i \setminus V_{i-1}$. By construction

$$\forall\, j < i; \quad ((x_1,\dots,x_n)(v_j\,s_j))_{v_i} \quad \neq \quad ((x_1,\dots,x_n)(v_i\,s_i))_{v_i}.$$

To prove (3) we can, without loss of generality, restrict ourselves to the case where we have an arbitrary permutation $(x_1,\dots,x_n)$ and $(y_1,\dots,y_n)$, the unique permutation satisfying $y_{v_i} = i$. We proceed by constructing a $\Gamma(S_n,T_n)$-path between these two permutations. Obviously, there exists an unique v_j such that $n = x_{v_j}$ and in the tree $\mathcal{T}_n$ there exists a unique path of length at most $\operatorname{diam}(\mathcal{T}_n) \le n-1$ connecting v_i and v_n. Accordingly, there is a $\Gamma(S_n,T_n)$-path of length at most $\operatorname{diam}(\mathcal{T}_n)$ between (x_i) and a permutation (z_i) such that $z_{v_n} = n$. Our construction in (2) implies

$$\forall\, i < n; \quad ((z_1,\dots,z_n)(v_i\,s_i))_{v_n} = n,$$

whence we can proceed inductively, moving $(n-1)$ to the v_{n-1}th position using the subtree $\mathcal{T}_{n-1}$. We consequently arrive at

$$\operatorname{diam}(\Gamma(S_n,T_n)) \le \sum_{i=2}^{n} \operatorname{diam}(\mathcal{T}_i) \le \binom{n}{2}$$

and the proof of the lemma is complete. □

In case of star-transpositions, i.e. $T_n = \{(1\,j) \mid 2 \le j \le n\}$, we have the following situation:

$$\{1\} \subset \{(1\,2)\} \subset \{(1\,2),(1\,3)\} \subset \cdots \subset \{(1\,j) \mid 2 \le j \le n\}, \tag{3.5}$$

$(v_i\, s_i) = (i+1\,1)$ i.e. $s_i = 1$ and $\mathrm{diam}(\Gamma(S_n, \{(1\,j) \mid 2 \le j \le n\})) \le 2(n-2)$. Similarly, in case of the bubble-sort graph, $T_n = \{(i\,i+1) \mid 1 \le i \le n-1\}$

$$\{1\} \subset \{(1\,2)\} \subset \{(1\,2),(2\,3)\} \subset \cdots \subset \{(i\,i+1) \mid 1 \le i \le n-1\} \tag{3.6}$$

we arrive at $(v_i\, s_i) = (i+1\,i)$ and $\mathrm{diam}(\Gamma(S_n, \{(i\,i+1) \mid 1 \le i \le n-1\})) = \binom{n}{2}$.

Lemma 2. *Suppose T_n is a minimal generating set of transpositions. We select permutations with independent probability $\lambda_n = \frac{1+\epsilon_n}{n-1}$, where $n^{-\frac{1}{3}+\delta} \le \epsilon_n$, for some $\delta > 0$. Then each permutation, v, is contained in a Γ_n-subtree $\mathcal{T}_n(v)$ of size $\lfloor \frac{1}{4} n^{\frac{2}{3}} \rfloor$ with probability at least $\wp(\epsilon_n)$.*

Proof. We construct the subtree $\mathcal{T}_n(v)$ by means of a branching process [15] within $\Gamma(S_n, T_n)$. Without loss of generality, we may initiate the process at id and assume (for notational convenience) $n - \frac{1}{2}n^{\frac{2}{3}} \in \mathbb{N}$. We shall begin by specifying an appropriate move-set (of transpositions) by which the offspring of the branching process is being generated. To this end, let

$$N = \{(v_j\, s_j) \mid 1 \le j \le n - \frac{1}{2}n^{\frac{2}{3}} - 1\} \subset T_n.$$

Note that N acts trivially on labels v_h where $h > n - \frac{1}{2}n^{\frac{2}{3}} - 1$.

The process is defined as follows: we set $U_0 = \varnothing \subset N$ and $M_0 = L_0 = \{id\} \subset S_n$. At step $(j+1)$, suppose we are given $U_j \subset N$, M_j and $L_j \subset S_n$. In case of $L_j = \varnothing$ or $|U_j| = \lfloor \frac{1}{4} n^{\frac{2}{3}} \rfloor - 1$ the process stops. Otherwise, we consider the smallest element $l_j \in L_j$ and select among its smallest $(n - \lfloor \frac{3}{4} n^{2/3} \rfloor - 1)$ neighbors, contained in $N \setminus U_j$ with independent probability λ_n. Let $x_1 = l_j\, r_{x_1}$ be the first selected l_j-neighbor and $r_{x_1} \in N \setminus U_j$. We then set $U_j(x_1) = U_j \dot\cup \{r_{x_1}\}$ and proceed the selection with the smallest $(n - \lfloor \frac{3}{4} n^{2/3} \rfloor - 1)$ neighbors contained in $N \setminus U_j(x_1)$ instead of those in $N \setminus U_j$. After all l_j neighbors are checked and given that $(x_1, \dots, x_s)$ have been subsequently selected, we set

$$\begin{aligned} U_{j+1} &= U_j \dot\cup \{r_{x_1}, \dots, r_{x_s}\} \\ L_{j+1} &= (L_j \setminus \{l_j\}) \cup \{x_1, \dots, x_s\} \\ M_{j+1} &= M_j \dot\cup \{x_1, \dots, x_s\}. \end{aligned}$$

The minimality of T_n and the fact that each T_n-element is used at most once implies that this process generates a tree, i.e. each M_{j+1}-element is considered only once. Furthermore, in view of

$$\frac{1+\epsilon_n}{n-1} \cdot \left(n - \lfloor \frac{3}{4} n^{\frac{2}{3}} \rfloor - 1\right) > 1 \tag{3.7}$$

this process can be viewed as a branching process embedded in $\Gamma(S_n, T_n)$. Therefore

$$\mathbb{P}\left(|M_j| = \lfloor \frac{1}{4} n^{\frac{2}{3}} \rfloor \mid \text{for some } j\right) \geq \wp(\epsilon_n)$$

and the lemma follows. $\square$

By choosing k sufficiently large, we next enlarge the Lemma 2-subtrees via an inductive construction to arbitrary polynomial size. We remark that Lemma 1 is of central importance for the construction of the Lemma 3-subtree, since some T_n-generators will be use multiple times.

Lemma 3. *Suppose k is arbitrary but fixed and let $\theta_n \geq O(n^{\delta})$. Then each Γ_n-vertex is contained in a Γ_n-subcomponent of size $\geq c_k n^{\frac{2}{3}} \theta_n^k \geq c_k n^{\frac{2}{3}+k\delta}$ with probability at least*

$$\delta_k(\epsilon_n) = \wp(\epsilon_n)\,(1 - e^{-\beta_k \theta_n}), \tag{3.8}$$

where $\beta_k > 0$ and $\epsilon_n \geq n^{-\frac{1}{3}+\delta}$.

Proof. Without loss of generality we may assume $\pi = id$, $\mu_n, \ell_n \in \mathbb{N}$ and set

$$A_m = \left\{(v_j^m\; s_j^m) \in T_n \mid 1 \leq j \leq \mu_n\right\}.$$

where $(v_j^m\; s_j^m) = (v_{z_n+j+(m-1)\mu_n-1}\; s_{z_n+j+(m-1)\mu_n-1})$. That is, A_m is the "first" (in the sense of the labeling given by the sequence $(v_2, v_3, \dots, v_n)$) subset of T_n-transpositions that act on labels v_i, where $i \leq z_n + m\mu_n - 1$ for $1 \leq m \leq k$.

We set $w_i^{(h)} = (v_j^h\; s_j^h)$ and consider the branching process of Lemma 2 at $\pi = id$, denoting the tree of size $\lfloor \frac{1}{4} n^{\frac{2}{3}} \rfloor$ by T^1. Let

$$Y_1 = \left|\{w_i^{(1)} \in A_1 \mid \exists x \in T^1; x \cdot w_i^{(1)} \in \Gamma_n\}\right|.$$

According to Lemma 1

$$\forall\, x, y \in T^1; \forall\, w_i^{(1)} \neq w_r^{(1)} \in A_1; \quad x \cdot w_i^{(1)} \neq y \cdot w_r^{(1)},$$

whence

$$\mathbb{E}[Y_1] = \mu_n \cdot \left(1 - \left(1 - \frac{1+\epsilon_n}{n-1}\right)^{\frac{1}{4}n^{\frac{2}{3}}}\right) \sim \mu_n \left(1 - \exp(-(1+\epsilon_n)\frac{1}{4} n^{-\frac{1}{3}})\right). \tag{3.9}$$

Using large deviation inequalities [9], we conclude that there exists some constant $c_1 > 0$ such that

$$\mathbb{P}\left(Y_1 < \frac{1}{2}\mathbb{E}[Y_1]\right) \leq \exp\left(-c_1 \cdot \mathbb{E}[Y_1]\right).$$

We select the smallest element, $x_{(i\,j)}$, from the set $\{x \cdot w_j^{(1)} \mid x \in T^1, x \cdot w_j^{(1)} \in \Gamma_n\}$ and start the branching process of Lemma 2 at $x_{(i\,j)}$. As a result, we derive the tree $C_2(x_{(i\,j)})$ of size $\lfloor \frac{1}{4} n^{\frac{2}{3}} \rfloor$ with probability at least $\wp(\epsilon_n)$. According to Lemma 2, the generation of this tree exclusively involves labels v_j where $j \leq z_n - 1$. Therefore, since any two smallest elements $x_{(i_1\,j_1)}$ and $x_{(i_2\,j_2)}$ differ in at least one of two coordinates with labels v_{j_1}, v_{j_2} for $z_n < j_1, j_2 \leq z_n + \mu_n$, we have

$$C_2(x_{(i_1\,j_1)}) \cap C_2(x_{(i_2\,j_2)}) = \varnothing.$$

Let X_1 be the r.v. counting the number of these new Γ_n-subcomponents. In view of eq. (3.9), we obtain

$$\mathbb{E}[X_1] \geq \wp(\epsilon_n) \cdot \mathbb{E}[Y_1] \sim \wp(\epsilon_n) \cdot \mu_n \left(1 - \exp(-(1+\epsilon_n)\frac{1}{4} n^{-\frac{1}{3}})\right) \triangleq \theta_n \geq O(n^{\delta}).$$

Again, using large deviation inequalities we conclude that there exists some $c_1 > 0$ such that

$$\mathbb{P}(X_1 < \frac{1}{2}\theta_n) \leq \exp(-c_1\theta_n)$$

or equivalently, since the union of all the $C_2(x_{(i\,j)})$-subcomponents with T^1 forms a $\Gamma(S_n, T_n)$-subtree, T^2, we have

$$\mathbb{P}\left(|T^2| < \lfloor \frac{1}{4} n^{2/3} \rfloor \cdot \frac{1}{2}\theta_n\right) \leq \exp(-c_1\theta_n). \tag{3.10}$$

We now proceed by induction:
Claim: For each $1 \leq i \leq k$ there exists some constant $c_i > 0$ and a $\Gamma(S_n, T_n)$-subtree T^i such that

$$\mathbb{P}(|T^i| < \lfloor \frac{1}{4} n^{2/3} \rfloor \cdot \frac{1}{2^i}\theta_n^i) \leq \exp(-c_i\theta_n).$$

We have already established the induction basis. As for the induction step, let us assume the claim holds for $i < k$ and let $C_i(\alpha)$ denote a subcomponent generated by the branching process of Lemma 2 in the i-th step. We consider the T_n-transpositions $w_r^{(i+1)} \neq w_a^{(i+1)} \in A_{i+1}$. We consider the minimal elements, x_r^{α} of

$$Y_{i+1} = \{w_r^{(i+1)} \in A_{i+1} \mid \exists x \in C_i(\alpha); x \cdot w_r^{(i+1)} \in \Gamma_n\}$$

at which we initiate the branching process of Lemma 2. The process generates tree subcomponents $C_{i+1}(x_r^{\alpha})$ of size $\lfloor \frac{1}{4} n^{\frac{2}{3}} \rfloor$ with probability $\geq \wp(\epsilon_n)$. Any two of these are mutually disjoint and let

X_{i+1} be the r.v. counting their number. We derive setting $q_n = \lfloor \frac{1}{4} n^{2/3} \rfloor$

$$\begin{aligned}
\mathbb{P}\left(|T^{i+1}| < q_n \frac{1}{2^{i+1}}\theta_n^{i+1}\right) &\le \underbrace{\mathbb{P}\left(|T^i| < q_n\frac{1}{2^i}\theta_n^i\right)}_{\text{failure at step } i} + \underbrace{\mathbb{P}\left(|T^{i+1}| < q_n \frac{1}{2^{i+1}}\theta_n^{i+1} \mid |T^i| \ge q_n \frac{1}{2^i}\theta_n^i\right)}_{\text{failure at step } i+1 \text{ conditional to } |T^i| \ge q_n \frac{1}{2^i}\theta_n^i} \\
&\le \underbrace{e^{-c_i\,\theta_n}}_{\text{induction hypothesis}} + \underbrace{e^{-c\,\theta_n^{i+1}}}_{\text{large deviation results}} \cdot (1 - e^{-c_i\,\theta_n}), \quad c_i > 0 \\
&\le e^{-c_{i+1}\,\theta_n}
\end{aligned}$$

and the claim follows.
Therefore each Γ_n-vertex is contained in a tree-subcomponent of size at least $O(n^{\frac{2}{3}}\theta_n^k) \ge O(n^{\frac{2}{3}+k\delta})$, with probability at least $\wp(\epsilon_n)(1-\exp(-c_k\theta_n))$ and the lemma is proved. $\square$

4. Vertices in small components

Let $\Gamma_{n,k}$ denote the set of Γ_n-vertices contained in components of size $\ge O(n^{k\delta+\frac{2}{3}})$ for some $0 < \delta < 1$. In this section we prove that $|\Gamma_{n,k}|$ is a.s. $\sim \wp(\epsilon_n)\frac{1+\epsilon_n}{n-1}n!$. In analogy to Lemma 3 of [30] we first observe that the number of vertices, contained in Γ_n-components of size $< c_k\, n^{k\delta+\frac{2}{3}}$, is sharply concentrated. The concentration reduces the problem to a computation of expectation values. It follows from considering the indicator r.vs. of pairs (C, v) where C is a component and $v \in C$ and to estimate their correlation. Since the components in question are small, no "critical" correlation terms arise.

Lemma 4. *Let* $\omega_n = |\Gamma_n \setminus \Gamma_{n,k}|$ *and* $\lambda_n = \frac{1+\epsilon_n}{n-1}$, *where* $n^{-\frac{1}{3}+\delta} \le \epsilon_n \le \lambda$, *for some* $\lambda > 0$. *Then we have*

$$\mathbb{P}\left(|\, \omega_n - \mathbb{E}[\omega_n]\, | \ge \frac{1}{n}\mathbb{E}[\omega_n]\right) = o(1). \tag{4.1}$$

With the help of Lemma 4, we proceed by computing the size of $\Gamma_{n,k}$.

Lemma 5. *Let* $\lambda_n = \frac{1+\epsilon_n}{n-1}$, *where* $n^{-\frac{1}{3}+\delta} \le \epsilon_n < 1$ *and suppose* $k \in \mathbb{N}$ *is sufficiently large. Then*

$$|\Gamma_{n,k}| \sim \wp(\epsilon_n)\frac{1+\epsilon_n}{n-1}n! \qquad a.s.\ . \tag{4.2}$$

Proof. First we prove for any $n^{-\frac{1}{3}+\delta} \le \epsilon_n \le \lambda$, where $\lambda > 0$

$$(1-o(1))\wp(\epsilon_n)\,|\Gamma_n| \le |\Gamma_{n,k}| \qquad \text{a.s.} \tag{4.3}$$

By assumption we have

$$\mathbb{E}[\omega_n] \le (1-\delta_k(\epsilon_n))|\Gamma_n|.$$

In view of Lemma 4, we derive

$$\omega_n < \left(1+O(\frac{1}{n})\right)\mathbb{E}[\omega_n] < \left(1-\delta_k(\epsilon_n)+O(\frac{1}{n})\right)|\Gamma_n| \quad \text{a.s.},$$

whence

$$|\Gamma_{n,k}| \ge \left(\delta_k(\epsilon_n)-O(\frac{1}{n})\right)|\Gamma_n| = (1-o(1))\wp(\epsilon_n)|\Gamma_n| \quad \text{a.s..}$$

Next we prove for $n^{-\frac{1}{3}+\delta} \le \epsilon_n < 1$

$$|\Gamma_{n,k}| \le (1+o(1))\wp(\epsilon_n)\,|\Gamma_n| \qquad \text{a.s.} \tag{4.4}$$

For this purpose we consider the branching process on a $(n-1)$-regular rooted tree T_{r^*} where the r.v. ξ_r^* of the rooted vertex r^* is $B(n-1,\lambda_n)$ distributed while the r.v. of any other vertex r has the distribution $B(n-2,\lambda_n)$. Let C_{r^*} denote the component generated by such a branching process. Bollobás *et al.* [8] showed that

$$\mathbb{P}(|C_{r^*}|=i) = (1+o(1))\cdot\frac{(\lambda_n\cdot(n-2))^{i-1}}{i\sqrt{2\pi i}}\left[\frac{(n-2)(1-\lambda_n)}{(n-3)}\right]^{ni-3i+2}, \tag{4.5}$$

where $i = i(n) \to \infty$ as $n \to \infty$. The key observation is an inequality due to Bollobás *et al.* [8], relating this process with the construction of a spanning tree of a Γ_n-component at vertex r,

$$\mathbb{P}\left(|C_{r^*}| \le m\right) \le \mathbb{P}\left(|C_r| \le m\right). \tag{4.6}$$

Eq. (4.6) follows immediately from the observation that during the generation of a spanning tree of a component, there are for each vertex *at most* $(n-2)$ neighbors, while in T_{r^*} there exist *exactly* $(n-2)$ neighbors. Suppose now k is sufficiently large, satisfying $k\delta+\frac{2}{3} \ge 3\frac{1}{3}$. Then $n^2 \le c_k\cdot n^{k\delta+\frac{2}{3}}$ for sufficiently large n, i.e.

$$\mathbb{P}(c_k\cdot n^{k\delta+\frac{2}{3}} \le |C_{r^*}| < \infty) \le \mathbb{P}(n^2 \le |C_{r^*}| < \infty). \tag{4.7}$$

This probability can be estimated as follows

$$\begin{aligned}\mathbb{P}(n^2 \le |C_{r^*}| < \infty) &= \sum_{i \ge n^2} \mathbb{P}(|C_{r^*}| = i) \\ &\le \sum_{i \ge n^2} (1+o(1)) \cdot \frac{(\lambda_n \cdot (n-2))^{i-1}}{i\sqrt{2\pi i}} \left[\frac{(n-2)(1-\lambda_n)}{(n-3)}\right]^{ni-3i+2} \\ &\le \sum_{i \ge n^2} \left[(1+\epsilon_n)e^{-\epsilon_n}\right]^i \le \sum_{i \ge n^2} c(\epsilon)^i = o(e^{-\ln(n)\, n}),\end{aligned}$$

where $0 < c(\epsilon) < 1$. We accordingly derive

$$\mathbb{P}(|C_{r^*}| < c_k \cdot n^{k\delta + \frac{2}{3}}) = \mathbb{P}(|C_{r^*}| < \infty) - \mathbb{P}(c_k \cdot n^{k\delta + \frac{2}{3}} \le |C_{r^*}| < \infty) \tag{4.8}$$

$$\ge (1 - \underbrace{(1+o(1))\wp(\epsilon_n)}_{=\pi_0}) - o(e^{-n^{\ln(n) n}}), \tag{4.9}$$

where π_0 denotes the survival probability of the branching process on T_{r^*}, see Corollary 1. From eq. (4.6) and eq. (4.9) we immediately obtain, taking the expectation

$$\mathbb{E}[\omega_n] \ge (1 - \wp(\epsilon_n))\, |\Gamma_n| + o(1). \tag{4.10}$$

Lemma 4 accordingly implies

$$(1 - \wp(\epsilon_n) - O(\frac{1}{n}))\, |\Gamma_n| \le \omega_n \qquad \text{a.s.}, \tag{4.11}$$

whence the lemma. □

We established in proof of Lemma 5 the relation

$$\mathbb{P}(n^2 \le |C_{r^*}| < \infty) \le \sum_{i \ge n^2} \left[(1+\epsilon_n)e^{-\epsilon_n}\right]^i \le \sum_{i \ge n^2} c(\epsilon)^i = o(e^{-\ln(n)\, n}).$$

Therefore, relaxing the condition $\epsilon_n < 1$, we loose the upper but not the lower bound. That is, we have for probabilities $\lambda_n = \frac{\lambda}{n-1}$ where $\lambda > 0$

Corollary 2. *Let $\lambda_n = \frac{\lambda}{n-1}$, where $0 < \lambda$ and suppose $k \in \mathbb{N}$ is sufficiently large. Then*

$$|\Gamma_{n,k}| \ge (1 - o(1))\, \pi_P(\lambda)\, \frac{\lambda}{n-1}\, n! \qquad a.s.\ . \tag{4.12}$$

5. The main theorem

We show in this section that the unique giant component forms within $\Gamma_{n,k}$ for two reasons: first, any $\Gamma_{n,k}$-vertex is *a priori* contained in a subcomponent of size $\geq O(n^{k\delta+\frac{2}{3}})$, limiting the number of ways by which $\Gamma_{n,k}$-splits can be chosen and second there are many independent paths connecting large $\Gamma(S_n,T_n)$-subsets. We first prove Lemma 6 according to which $\Gamma_{n,k}$ is "almost" 2-dense in $\Gamma(S_n,T_n)$.

Lemma 6. *Let* $k \in \mathbb{N}$*,* $\lambda_n = \frac{1+\epsilon_n}{n-1}$ *where* $\epsilon_n \geq n^{-\frac{1}{3}+\delta}$ *for some* $\delta > 0$ *and let furthermore* $A_\delta = \left\{v \mid |d(v,2)\cap\Gamma_{n,k}| < c_k \cdot n^\delta\right\}$*. Then we have for some* $\kappa > 0$*,* $\mathbb{P}(v \in A_\delta) \leq \exp(-\kappa n^\delta)$ *and there exists some* $0 < \rho < \kappa$ *such that*

$$|A_\delta| \leq n!\exp(-\rho n^\delta) \quad a.s..$$

Proof. We consider now the action of the transpositions

$$A_{k+1} = \left\{(v_j^{k+1}\ s_j^{k+1}) \in T_n \mid 1 \leq j \leq \ell_n\right\}$$

where $w_j^{(k+1)} = (v_j^{k+1}\ s_j^{k+1}) = (v_{z_n-1+j+k\mu_n}\ s_{z_n-1+j+k\mu_n})$ and set

$$d^{(k+1)}(v,2) = \{v \cdot w_i^{(k+1)} \cdot w_j^{(k+1)} | 1 \leq i < j \leq \ell_n\}.$$

We proceed by establishing a lower bound on the cardinality of $d^{(k+1)}(v,2)$. Since T_n is a minimal generating set, any sequence of distinct T_n-transpositions is acyclic. Therefore

$$|d^{(k+1)}(v,2)| \quad \geq \quad \frac{1}{2!}\binom{\ell_n}{2} = \frac{n^{\frac{4}{3}}}{(2!)^2}\cdot\left[\frac{k}{2(k+1)}\right]^2\cdot(1-o(1)).$$

Let $\Delta_k = \left[\frac{k}{2(k+1)}\right]^2/(2!)^2$ and Z be the r.v. counting the number of vertices contained in the set $d^{(k+1)}(v,2)\cap\Gamma_{n,k}$, whose subcomponents are the trees constructed in Lemma 3. We immediately compute

$$\mathbb{E}(Z) \geq \lambda_n \cdot \delta_k(\epsilon_n)\cdot|d^{(k+1)}(v,2)| \sim \Delta_k\, n^{\frac{4}{3}}\cdot\frac{1+\epsilon_n}{n-1}\cdot\wp(\epsilon_n)(1-e^{-\beta_k\theta_n}) \geq \Delta_k \cdot n^\delta.$$

The key observation is the following: the construction of the Lemma 3-subtrees did not involve any labels $v_{z_n-1+j+k\mu_n}$, i.e. any two such subtrees remain vertex-disjoint. Therefore the r.v. Z is a sum of *independent* indicator r.vs. and Chernoff's large deviation inequality [9] implies

$$\mathbb{P}(v \in A_\delta) \leq \mathbb{P}\left(Z < \frac{1}{2}\Delta_k\cdot n^\delta\right) \leq \exp(-\kappa n^\delta) \quad \text{for some } \kappa > 0. \tag{5.1}$$

Consequently, the expected number of vertices contained in A_δ is bounded by $n!\exp(-\kappa n^\delta)$. Now Markov's inequality $\mathbb{P}(X > t\mathbb{E}(X)) \le 1/t$, for $t > 0$, guarantees $|A_\delta| \le n! \cdot e^{-\rho n^\delta}$ a.s. for any $0 < \rho < \kappa$ and the lemma follows. □

Next we show that there exist many vertex disjoint paths between $\Gamma_{n,k}$-splits of sufficiently large size. The proof is analogous to Lemma 7 in [30]. We remark that Lemma 7 does not use an isoperimetric inequality [14]. It only employs a generic estimate of the vertex boundary in Cayley graphs due to Aldous [4, 6].

Lemma 7. *Let (S,T) be a vertex-split of $\Gamma_{n,k}$ with the properties*

$$\exists\, 0 < \rho_0 \le \rho_1 < 1; \quad (n-2)! \le |S| = \rho_0|\Gamma_{n,k}| \quad \text{and} \quad (n-2)! \le |T| = \rho_1|\Gamma_{n,k}|. \tag{5.2}$$

Then there exists some $c > 0$ such that a.s. $d(S)$ is connected to $d(T)$ in $\Gamma(S_n, T_n)$ via at least

$$c\,(n-5)!/(n-1)^7 \tag{5.3}$$

vertex disjoint (independent) paths of length ≤ 3.

Proof. We distinguish the cases $|\mathsf{B}(S,2)| \le \frac{2}{3}\,n!$ and $|\mathsf{B}(S,2)| > \frac{2}{3}\,n!$. In the former case, we employ the generic estimate of vertex boundaries in Cayley graphs [4]

$$|\mathsf{d}(S)| \ge \frac{1}{\mathrm{diam}(\Gamma(S_n,T_n))} \cdot |S| \left(1 - \frac{|S|}{n!}\right). \tag{5.4}$$

In view of eq. (5.2) and Lemma 1, eq. (5.4) implies

$$\exists\, d_1 > 0; \quad |\mathsf{d}(\mathsf{B}(S,2))| \ge \frac{d_1}{n^2} \cdot |\mathsf{B}(S,2)| \ge d_1 \cdot (n-4)!. \tag{5.5}$$

According to Lemma 6, a.s. all but $\le n!\,e^{-\tilde{\Delta} n^\delta}$ permutations are within distance 2 to some $\Gamma_{n,k}$-vertex, whence

$$|\mathsf{d}(\mathsf{B}(S,2)) \cap \mathsf{B}(T,2)| \ge d_2 \cdot (n-4)! \quad \text{a.s..} \tag{5.6}$$

Let $\beta_2 \in \mathsf{d}(\mathsf{B}(S,2)) \cap \mathsf{B}(T,2)$. Then there exists a path $(\alpha_1, \alpha_2, \beta_2)$ such that $\alpha_1 \in \mathsf{d}(S)$, $\alpha_2 \in \mathsf{d}(B(S,1))$. We distinguish the cases

$$|\mathsf{d}(\mathsf{B}(S,2)) \cap \mathsf{d}(\mathsf{B}(T,1))| \ge d_{2,1}\,(n-4)! \quad \text{and} \quad |\mathsf{d}(\mathsf{B}(S,2)) \cap \mathsf{B}(T,1)| \ge d_{2,2}\,(n-4)!. \tag{5.7}$$

For $|\mathsf{d}(\mathsf{B}(S,2)) \cap \mathsf{d}(\mathsf{B}(T,1))| \ge d_{2,1}\,(n-4)!$, we consider the set

$$T^* = \{\beta_1 \in \mathsf{d}(T) \mid d(\beta_1, \beta_2) = 1, \text{for some } \beta_2 \in \mathsf{d}(\mathsf{B}(T,1))\}.$$

Evidently, at most $n-1$ elements in $\mathsf{d}(T)$ can be connected to a fixed β_2, whence

$$|T^*| \geq \frac{1}{2} d_{2,1}\,(n-5)!.$$

Let $T_1 \subset T^*$ be some maximal set such that any pair of T_1-vertices (β_1, β_1') has at least distance $d(\beta_1, \beta_1') > 6$. Then $|T_1| > |T^*|/(n-1)^7$ since $|\mathsf{B}(v,7)| < (n-1)^7$. Any two of the paths from $\mathsf{d}(S)$ to $T_1 \subset \mathsf{d}(T)$ are of the form $(\alpha_1, \alpha_2, \beta_2, \beta_1)$ and vertex disjoint since each of them is contained in $\mathsf{T}(\beta_1, 3)$. Accordingly there are a.s. at least

$$\frac{1}{2} d_{2,1}\,(n-5)!/(n-1)^7 \tag{5.8}$$

vertex disjoint paths connecting $\mathsf{d}(S)$ and $\mathsf{d}(T)$. In case of $|\mathsf{d}(\mathsf{B}(S,2)) \cap \mathsf{B}(T,1)| \geq d_{2,2}\,(n-3)!$ we analogously conclude, that there exist a.s. at least

$$d_{2,2}\,(n-4)!/(n-1)^5 \tag{5.9}$$

vertex disjoint paths of the form $(\alpha_1, \alpha_2, \beta_2)$ connecting $\mathsf{d}(S)$ and $\mathsf{d}(T)$.
It remains to consider the case $|\mathsf{B}(S,2)| > \frac{2}{3} \cdot n!$. By construction both: S and T satisfy eq. (5.2), whence we can, without loss of generality assume that also $|\mathsf{B}(S,2)| > \frac{2}{3} \cdot n!$ holds. But then

$$|\mathsf{B}(S,2) \cap \mathsf{B}(T,2)| > \frac{1}{3}\, n!$$

and for each $\alpha_2 \in \mathsf{B}(S,2) \cap \mathsf{B}(T,2)$ we select $\alpha_1 \in \mathsf{d}(S)$ and $\beta_1 \in \mathsf{d}(T)$. We derive in analogy to the previous arguments that there exist a.s. at least

$$d_2\,(n-3)!/(n-1)^5 \tag{5.10}$$

pairwise vertex disjoint paths of the form $(\alpha_1, \alpha_2, \beta_1)$ and the proof of the lemma is complete. $\square$

Proof of Theorem 1. To prove the theorem we employ an argument due to Ajtai *et al.* [1] originally used for n-cubes and independent edge-selection. We proceed along the lines of [30] and select the $\Gamma(S_n, T_n)$-vertices in two distinct randomizations.
Let $x_1, x_2 > 0$ such that $\frac{1}{x_1} + \frac{1}{x_2} = 1$. First we select with probability $\frac{1+\epsilon_n/x_1}{n}$ and second with probability $\frac{\epsilon_n}{x_2 \cdot n}$. The probability of not being chosen in both rounds is given by

$$\left(1 - \frac{1+\epsilon_n/x_1}{n}\right)\left(1 - \frac{\epsilon_n}{x_2 \cdot n}\right) \geq 1 - \frac{1+\epsilon_n}{n},$$

whence it suffices to prove that after the second randomization there exists a giant component with the property $|C_n^{(1)}| \sim |\Gamma_{n,k}|$.

After the first randomization each $\Gamma(S_n,T_n)$-vertex has been selected with probability $\frac{1+\epsilon_n/x_1}{n}$ and according to Lemma 5, we have

$$|\Gamma_{n,k}(x_1)| \sim \wp(\epsilon_n/x_1)\,|\Gamma_n(x_1)| \quad \text{a.s.,} \tag{5.11}$$

where $\Gamma_n(x_1)\subset\Gamma_n$. Suppose $\Gamma_{n,k}(x_1)$ contains a "large" component, S. To be precise a component S of size

$$(n-2)! \le |S| \le (1-b)\,|\Gamma_{n,k}(x_1)|, \quad \text{where } b>0.$$

Then there exists a split of $\Gamma_{n,k}(x_1)$, (S,T), satisfying the assumptions of Lemma 7 (and, of course, $d(S)\cap d(T)=\varnothing$). We observe that Lemma 3 limits the number of ways these splits can be constructed. Let $M_k \triangleq \lfloor \frac{1}{4}n^{\frac{2}{3}}\rfloor \theta_n^k \ge c_k\, n^{k\delta+\frac{2}{3}}$ for some $c_k>0$. Obviously, there are at most $2^{n!/M_k}$ ways to select S of such a split. Now we employ Lemma 7. In view of $(n-2)!\le |S|$, Lemma 7 implies that there exists some $c>0$ such that a.s. $d(S)$ is connected to $d(T)$ in $\Gamma(S_n,T_n)$ via at least $c\cdot n!/n^{12}\le c\cdot |S|/n^{10}$ vertex disjoint paths of length ≤ 3.
We next perform the second randomization and select $\Gamma(S_n,T_n)$-vertices with probability $\frac{\epsilon_n/x_2}{n}$. None of the above $c\cdot|S|/n^{10}$ paths can be selected during this process. Since any two paths are vertex disjoint the expected number of such splits is, by linearity of expectation, less than

$$2^{n!/M_k}(1-(\epsilon_n/x_2 n)^4)^{\frac{c\cdot n!}{n^{12}}} \le 2^{n!/M_k}e^{-c' n!/n^{16}} \quad \text{for some } c,c'>0. \tag{5.12}$$

Accordingly, choosing k sufficiently large the expected number of these $\Gamma_{n,k}(x_1)$-splits tends to zero, i.e. for any $k\ge k_0\in\mathbb{N}$ there exists a.s. no two component split (S,T) of $\Gamma_{n,k}(x_1)$ with the property $\rho_0|\Gamma_{n,k}(x_1)| = |S| \le |T|$. Consequently, there exists some subcomponent $C_n(x_1)$ with the property

$$|C_n(x_1)| = |\Gamma_{n,k}(x_1)| \sim \wp(\epsilon_n/x_1)\,|\Gamma(x_1)| \quad \text{a.s.,}$$

obtained by the merging of the subcomponents of size $\ge O(n^{k\delta+\frac{2}{3}})$ generated during the first randomization via the paths selected during the second. Since $\wp(\epsilon_n/x_1)$ is continuous in the parameter ϵ_n/x_1, see eq. (3.1), we derive, for x_1 tending to 1

$$|C_n^{(1)}| = \lim_{x_1\to 1} |C_n(x_1)| \sim \wp(\epsilon_n)|\Gamma_n| \quad \text{a.s.} \tag{5.13}$$

It remains to prove uniqueness. Any other largest component, $\tilde{C}_n$, is necessarily contained in $\Gamma_{n,k}$. However, we have just proved $|C_n^{(1)}|\sim\wp(\epsilon_n)|\Gamma_n|$ and according to Lemma 5, $\wp(\epsilon_n)|\Gamma_n|\sim|\Gamma_{n,k}|$. Therefore $|\tilde{C}_n| = o(|C_n^{(1)}|)$, whence $C_n^{(1)}$ is unique. □

Acknowledgments.

We are grateful to Fenix W.D. Huang and Rita R. Wang for their help. This work was supported by the 973 Project, the PCSIRT Project of the Ministry of Education, the Ministry of Science and Technology, and the National Science Foundation of China.

References

[1] M. Ajtai, J. Komlós and E. Szemerédi, Largest random component of a k-cube, *Combinatorica* **2**(1982), 1-7.

[2] S.B. Akers, D. Harel and B. Krishnamurthy, The star graph: An attractive alternative to the n-cube, *Proceedings of the International Conference on Parallel Processing* 1987, 393-400.

[3] S.B. Akers and B. Krishnamurthy, A group theoretic model for symmetric interconnection networks, *IEEE Transactionsaction on Computers* **38**(1989), 555-565.

[4] D. Aldous and P. Diaconis, Strong uniform times and finite random walks, *Adv. in Appl. Math.* **2**(1987), 69-97.

[5] T. Araki, Hyper hamiltonian laceability of Cayley graphs generated by transpositions, *Networks* 2006, 121-124.

[6] L. Babai, Local expansion of vertex transitive graphs and random generation in finite groups, *Proc* 23 *ACM Symposium on Theory of Computing (ACM New York)* **1**(1991), 164-174.

[7] V. Bafna and P. A. Pevzner, Sorting by Reversals: Genome Rearrangements in Plant Organelles and Evolutionary History of X Chromosome, *Mol. Biol. and Evol.* 1995.

[8] B. Bollobás, Y. Kohayakawa and T. Luczak, The evolution of random subgraphs of the cube, *Random Struct. Alg.* **3**(1992), 55-90.

[9] H. Chernoff. A measure of the asymptotic efficiency for tests of a hypothesis based on the sum of observations, *Annals of Mathematical Statistics* **23**(1952), 493-509.

[10] B.S. Chlebus, K. Diks and A. Pelc, Reliable Broadcasting in Hypercubes with Random Link and Node Failures, *Combinatorics, Probability and Computing* **5**(1996), 337-350.

[11] N. Eriksen and A. Hultman, Estimating the expected reversal distance after a fixed number of reversals, *Advances in Appl. Math.* **32**(2004), 439-453.

[12] A.H. Esfahanian, Generalized Measures of Fault Tolerance with Application to n-Cube Networks, *IEEE Transactions on Computers* **38**(11)(1993), 1586-1591.

[13] L.J. Fan, C.B. Yang and S.H. Shiau, Routing algorithms on the bus-based hypercube network, *IEEE Transactions on Parallel and Distributed Systems* **16**(4)(2005), 335-348.

[14] L.H. Harper, Minimal numberings and isoperimetric problems on cubes, *Theory of Graphs, International Symposium, Rome* 1966.

[15] T.E. Harris, The Theory of Branching Processes (Dover Phenix editions), Dover Pubns. *Springer Verlag* 1963.

[16] S.Y. Hsieh, G.H. Chen and C.W. Ho, Hamiltonian-laceability of star graphs, *Networks* **36**(2000), 225-232.

[17] S.Y. Hsieh, Embedding longest fault-free paths onto star graphs with more vertex faults, *Theoretical Computer Science* **337**(2005), 370-378.

[18] S.Y. Hsieh, G.H. Chen and C.W. Ho, Longest fault-free paths in star graphs with vertex faults, *Theoretical Computer Science* **262**(2001), 215-227.

[19] S.Y. Hsieh, G.H. Chen and C.W. Ho, Longest fault-free paths in star graphs with edge faults, *IEEE Transactions on Computers* **50**(2001), 960-971.

[20] J. Irving and A. Rattan, Minimal Factorizations of Permutations into Star Transpositions, *FPSAC* 2008.

[21] J.S. Jwo, S. Lakshmivarahan and S.K. Dhall, Embedding of cycles and grids in star graphs, *Journal of Circuits, Systems, and Computers* **1**(1991), 43–74.

[22] N. Khiripet, Bacterial whole genome phylogeny using proteome comparison and optimal reversal distance, *Computational Systems Bioinformatics Conference, IEEE.* (2005).

[23] V.F. Kolchin, Random Mappings, Optimization Software Inc., *Springer Verlag* New York, 1986.

[24] S. Latifi, On the fault-diameter of the star graph, *Information Processing Letters* **46**(1993), 143-150.

[25] T.K. Li, J.J.M. Tan and L.H. Hsu, Hyper hamiltonian laceability on edge fault star graph, *Information Sciences* **165**(2004), 59-71.

[26] C.K. Lin, J.J.M. Tan, L.H. Hsu, E. Cheng, L. Liptak, Conditional Diagnosability of Cayley Graphs Generated by Transposition Trees under the Comparison Diagnosis Model, *Journal of Interconnection Networks* **9**(1-2)(2008).

[27] K. Padmanabhan, The composite binary cubea family of interconnection networks for multiprocessors, *Proceedings of the* 3*rd international conference on Supercomputing* 1989, 62-71.

[28] I. Pak, Reduced decompositions of permutations in terms of star transpositions, generalized catalan numbers and k-ary trees, *Discr. Math.* **204**(1999), 329-335.

[29] S. Hannenhalli and P.A. Pevzner, Transforming cabbage into Turnip (Polynomial Algorithm for sorting signed permutations by reversals), *Journal of the ACM* **48**(1999), 1-27.

[30] C. Reidys, Large components in random induced subgraphs of n-cubes, *Discr. Maths.* **309**(10)(2009), 3113-3124.

[31] Y. Rouskov, S. Latifi and P.K. Srimani, Conditional fault diameter of star graph networks, *Journal of Parallel and Distributed Computing* **33**(1996), 91-97.

[32] V. Sharma and E.M. Varvarigos, Some closed form results for circuit switching in a hypercube network, *Lecture Notes in Computer Science* 2006, 1124-1996.

[33] M. Tchuente, Generation of permutations by graphical exchanges, *Ars Combinatoria* **14**(1982), 115-122.

Center for Combinatorics, LPMC-TJKLC, Nankai University, Tianjin 300071, P.R. China, Phone: *86-22-2350-6800, Fax: *86-22-2350-9272

E-mail address: `duck@santafe.edu`